

DOI: <https://doi.org/10.22141/2307-1257.12.4.2023.424>

EULAR 2023 recommendations for SLE treatment:
synopsis for the management of lupus nephritis —
the European Renal Association (ERA) —
Immunonephrology Working Group (ERA-IWG)
perspective

Eleni Frangou¹, Annette Bruchfeld^{2,3}, Gema M. Fernandez-Juarez⁴, Jürgen Floege⁵,
Dimitrios Goumenos⁶, Sarah M. Moran⁷, Stefanie Steiger⁸, Kate I. Stevens⁹, Kultigin Turkmen¹⁰,
Andreas Kronbichler¹¹

¹Department of Nephrology, Limassol General Hospital, Limassol, Cyprus; University of Nicosia Medical School, Nicosia, Cyprus

²Department of Health, Medicine and Caring Sciences, Linköping University, Linköping, Sweden

³Department of Renal Medicine, Karolinska University Hospital and CLINTEC Karolinska Institutet, Stockholm, Sweden

⁴Department of Nephrology, Hospital Universitario La Paz, Madrid, Spain; Instituto de Investigacion la Paz, IDIPaZ, Spain

⁵Division of Nephrology, RWTH Aachen University Hospital, Aachen, Germany

⁶Department of Nephrology and Renal Transplantation, Patras University Hospital, Patras, Greece

⁷Cork University Hospital, University College Cork, Cork, Ireland

⁸Renal Division, Department of Medicine IV, Ludwig-Maximilians-University Hospital Munich, Ludwig-Maximilians-University Munich, Germany

⁹Glasgow Renal and Transplant Unit, Queen Elizabeth University Hospital, Glasgow, UK

¹⁰Division of Nephrology, Department of Internal Medicine, Necmettin Erbakan University, Konya, Turkey

¹¹Department of Internal Medicine IV, Nephrology and Hypertension, Medical University Innsbruck, Innsbruck, Austria




Figure 1. Specific considerations when evaluating the optimal currently approved add-on therapy of a patient with active lupus nephritis (LN) class III–V

Notes: BEL — belimumab; eGFR — estimated glomerular filtration rate; i.v. — intravenous; s.c. — subcutaneous; VCS — voclosporin.